Submit your solutions in writing no later than May 20 or present them orally during my office hours (MWF 3–4 in 19D).

1. Suppose A is a locally compact Hausdorff abelian group (typical cases include $A = \mathbf{R}$, $A = \mathbf{Z}$, and $A = \mathrm{U}(1)$). Recall (Example 7.18) the Pontryagin dual group $\mathrm{PD}(A) = \mathrm{Hom}(A, \mathrm{U}(1))$. Recall also the two functors of the form LocCompHausAb $\rightarrow \mathrm{Ban}_1$, namely, L¹ \circ HaarMeas and C₀ \circ PD. Here L¹(HaarMeas(A)) (henceforth L¹(A)) is the Banach space of finite measures on the measurable space HaarMeas(A) and C₀(PD(A)) is the Banach space of **C**-valued continuous functions on the topological group PD(A) that vanish at infinity. The *Fourier transform* on A is defined as a bounded map $F_A: \mathrm{L}^1(A) \rightarrow \mathrm{C}_0(\mathrm{PD}(A))$ that sends $\mu \in \mathrm{L}^1(A)$ to the function $\chi \mapsto \int \chi \cdot \mu$, where $\chi \in \mathrm{PD}(A)$, i.e., $\chi: A \to \mathrm{U}(1)$, and $\chi \cdot \mu$ denotes the product of a measurable function and a finite measure. Prove that the collection of maps F_A defined above is a natural transformation from L¹ \circ HaarMeas to C₀ \circ PD.

2. Prove that the category Ban_1 admits arbitrary small coproducts. Prove that the category Ban admits finite coproducts, but does not admit infinite coproducts.

3. Suppose V and W are two Banach spaces in the category Ban_1 of Banach spaces and contractive linear maps. A contractive bilinear map is a bilinear map $b: V, W \to A$ such that $||b(v, w)|| \leq 1$ for all $v \in V$ and $w \in W$ such that $||v|| \leq 1$ and $||w|| \leq 1$. Prove the existence of a Banach space $V \otimes_{\mathbf{C}} W$ with the following universal property: contractive bilinear maps $V, W \to A$ are in natural bijection with contractive linear maps $V \otimes_{\mathbf{C}} W \to A$. Prove that the span of elements of the form $v \otimes_{\mathbf{C}} w$ is dense in $V \otimes_{\mathbf{C}} W$. Does it always coincide with $V \otimes_{\mathbf{C}} W$?

4. Suppose S is a set and $P \subset S \times S$ is a set of pairs of elements in S. Prove that the quotient set S/\sim , where \sim is the equivalence relation generated by P, is isomorphic to the coequalizer of $P \xrightarrow{p_0}_{p_1} S$, where $p_0, p_1: P \to S$ are the projection maps.

5. Denote by $U: \mathsf{CAlg}_k \to \mathsf{Vect}_k$ the forgetful functor from the category of commutative k-algebras to the category of vector spaces over k. Fix a vector space $V \in \mathsf{Vect}_k$. Consider the functor $F: \mathsf{CAlg}_k \to \mathsf{Set}$ such that F(A) is the set of linear maps $V \to U(A)$ and for a homomorphism $f: A \to A'$ the map F(f) sends $g: V \to U(A)$ to $f \circ g: V \to U(A')$. Prove that F is corepresentable. Identify its corepresenting object as a familiar object from your algebra course.

6. Fix some vector spaces $V, W \in \mathsf{Vect}_k$. Consider the functor $F: \mathsf{Vect}_k^{\mathsf{op}} \to \mathsf{Set}$ such that F(A) is the set of linear maps $A \otimes_k V \to W$ and F(f) for a linear map $f: A \to A'$ sends a map $A' \otimes_k V \to W$ to its composition with $A \otimes_k V \to A' \otimes_k V$. Prove that F is representable. Identify its representing object as a familiar object from your algebra course. Same question when Vect_k is replaced by Ban_1 and \otimes_k is replaced by $\hat{\otimes}_{\mathbf{C}}$.