

## L<sup>p</sup>-SPACES ASSOCIATED WITH AN ARBITRARY VON NEUMANN ALGEBRA

by

UFFE HAAGERUP

Odense Universitet, Denmark.

Abstract: To any von Neumann algebra  $M$ , we associate Banach spaces  $L^p(M)$ ,  $1 \leq p \leq \infty$ , which generalize the classical Banach spaces  $L^p(\Omega, \mu)$  of functions on a measure space  $(\Omega, \mu)$ . We show that  $L^\infty(M) \cong M$ ,  $L^1(M) \cong M_*$ , and that  $L^2(M)$  is isomorphic to the Hilbert space of  $M$  in its standard form. When  $M$  is semifinite, the  $L^p(M)$ -spaces are isometric isomorphic to the spaces  $L^p(M, \tau)$  introduced by Dixmier, Segal and Kunze in 1953-1958. The  $L^p(M)$ -spaces are constructed as certain spaces of unbounded operators affiliated with the crossed product  $R(M, \sigma^\varphi)$  of  $M$  with the modular automorphism group associated with a fixed weight  $\varphi$  on  $M$ . The construction turns out to be independent (up to unitary equivalence) of the choice of  $\varphi$ .

**RESUMÉ** A toute algèbre de Von Neumann  $M$  nous associons des espaces de Banach  $L^p(M)$ ,  $1 \leq p \leq \infty$ , qui généralisent les espaces de Banach classiques  $L^p(\Omega, \mu)$  de fonctions sur un espace mesuré  $(\Omega, \mu)$ . Nous montrons que  $L^\infty(M) \cong M$ ,  $L^1(M) \cong M_*$ , et que  $L^2(M)$  est isomorphe à l'espace de Hilbert de la représentation standard. Si  $M$  est semi-finie les espaces  $L^p(M)$  sont isométriquement isomorphes aux espaces  $L^p(M, \tau)$  introduits par Dixmier, Segal et Kunze en 1953-1958. Les espaces  $L^p(M)$  sont construits comme espaces d'opérateurs non bornés affiliés aux produits croisés  $R(M, \sigma^\varphi)$  de  $M$  avec l'automorphisme modulaire associé à un poids fixe  $\varphi$  sur  $M$ . La construction s'avère indépendante (à une équivalence unitaire près) du choix de  $\varphi$ .

## Introduction

This note contains an outline of a forthcoming paper.

In [4], [11] and [8] J. Dixmier, I. Segal and R. Kunze have constructed the  $L^p$ -spaces  $L^p(M, \tau)$  associated with a semifinite von Neumann algebra  $M$ , which generalize the classical Banach spaces  $L^p(\Omega, \mu)$ . The  $L^p$ -spaces we construct in this note will consist of operators affiliated not with  $M$  itself but with a bigger algebra, namely the crossed product  $M_0 = R(M, \sigma^\varphi)$  of  $M$  with a modular automorphism group.  $M_0$  has a trace  $\tau$  satisfying  $\tau \circ \theta_s^\varphi = e^{-s} \tau$  where  $\theta_s^\varphi$  is the dual action.  $L^p(M)$  is defined as the set of  $\tau$ -measurable operators  $h$  affiliated with  $M_0$  satisfying

$$\begin{cases} \theta_s^\varphi h = \exp(-\frac{s}{p})h & p < \infty \\ \theta_s^\varphi h = h & p = \infty \end{cases}$$

equipped with a suitable norm. Since the triple  $(M_0, \tau, \theta^\varphi)$  is independent (up to unitary equivalence) of the choice of  $\varphi$ , the  $L^p$ -spaces are independent of  $\varphi$ .

We have  $L^\infty(M) = M$  and  $L^1(M) \simeq M_*$ .  $L^2(M)$  is a Hilbert space, and the representation of  $L^\infty(M)$  on  $L^2(M)$  defined by left multiplication is standard. If  $M$  is semifinite the  $L^p$ -spaces constructed in this way are isometric- and orderisomorphic to  $L^p(M, \tau_0)$  for any n.f.s. trace  $\tau_0$  on  $M$ .

## § 1 Construction of the $L^p$ -spaces

Let  $M$  be a von Neumann algebra. We will identify  $M$  with its natural injection in the crossed product  $M_0 = R(M, \sigma^{\varphi_0})$  where  $\varphi_0$  is a fixed weight on  $M$ . By construction  $M_0$  is generated by  $M$  and a one parameter group of unitaries  $\lambda(t)$  such that for  $x \in M$ ,  $\sigma_t^{\varphi_0}(x) = \lambda(t)x\lambda(t)^*$ .

Let  $T$  be the operator valued weight,  $T: M_0^+ \rightarrow \hat{M}_+$ , given by

$$T(x) = \int_{\hat{R}} \theta_s(x) ds, \quad x \in M_0^+,$$

where  $\theta_s = \sigma_s^{\varphi_0}$  is the dual action on  $M_0$  (cf [6], [7]). The weight  $\varphi_0 \circ T$  on  $M_0$  is  $2\pi$  times the dual weight to  $\varphi_0$ . Hence there exists a trace  $\tau$  on  $M_0$ , such that  $\varphi_0 \circ T = \tau(h \cdot)$  where  $h$  is the positive selfadjoint operator affiliated with  $M_0$  determined by  $h^{it} = \lambda(t)$  (cf [12], proof of lemma 8.2). The trace  $\tau$  satisfies

$$\tau \circ \theta_s = e^{-s} \tau, \quad s \in \hat{R}$$

### 1.1 Definition

For any normal semifinite weight  $\varphi$  on  $M$  we put  $\tilde{\varphi} = \varphi \circ T$ , and we let  $h_\varphi$  be the Radon-Nikodym derivative of  $\tilde{\varphi}$  with respect to the trace  $\tau$  on  $M_0$ , i.e.  $\tilde{\varphi} = \tau(h_\varphi \cdot)$ .

### 1.2 Theorem

1) The set  $\{h_\varphi \mid \varphi \text{ normal, semifinite}\}$  is equal to the set of positive selfadjoint operators  $h$  affiliated with  $M_0$ , which satisfies

$$\theta_s h = e^{-s} h \quad s \in \hat{R}$$

2) If  $\int_0^\infty \lambda de_\lambda^{\varphi}$  is the spectral decomposition of  $h_\varphi$  then

$$\tau((e_\lambda^{\varphi})^\perp) = \frac{1}{\lambda} \varphi(1), \quad \lambda > 0.$$

In particular  $h_\varphi$  is  $\tau$ -measurable iff  $\varphi$  is bounded (cf [9] p. 111).

### 1.3 Theorem

The map  $\varphi \rightarrow h_\varphi$ ,  $\varphi \in M_*^+$  has a unique extension to a linear map of  $M_*$  onto the set of  $\tau$ -measurable operators  $h$  affiliated with  $M_0$ , satisfying

$$\theta_s h = e^{-s} h.$$

(Note that the set of  $\tau$ -measurable operators on  $M_0$  is an algebra with respect to strong sum and strong product, cf [11]).

#### 1.4 Definition

- 1) We let  $L^1(M)$  denote the set of  $\tau$ -measurable operators  $h$  affiliated with  $M_0$ , for which  $\theta_s h = e^{-s} h$ ,  $s \in \hat{\mathbb{R}}$ .
- ii) We define a linear functional  $\text{tr}$  on  $L^1(M)$  by  $\text{tr}(h_\varphi) = \varphi(1)$ .

#### 1.5 Proposition

The map  $\varphi \rightarrow h_\varphi$  of  $M_*$  onto  $L^1(M)$  is an isometry with respect to the norm  $\|h\|_1 = \text{tr}(|h|)$  on  $L^1(M)$ .

#### 1.6 Remarks

- a) The trace  $\tau$  is infinite on any non vanishing operator in  $L^1(M)$ .
- b) If  $M$  is a factor of type III<sub>1</sub>,  $M_0$  is a II<sub>∞</sub>-factor. In this case any normal trace on  $M_0$  is proportional to  $\tau$ . Hence  $\text{tr}$  is not in general the restriction of a trace on  $M_0$ .

#### 1.7 Definition

We put  $L^p(M) = \{h, \tau\text{-measurable aff. with } M_0 \mid \theta_s h = \exp(-\frac{s}{p})h\}$   
and  $L^\infty(M) = \{h, \tau\text{-measurable aff. with } M_0 \mid \theta_s h = h\}$

#### 1.8 Remarks

- a) If  $p \neq q$  then  $L^p(M) \cap L^q(M) = \{0\}$ .
- b) If  $p < \infty$  then any non vanishing  $L^p$ -operator is unbounded.
- c)  $L^\infty(M)$  consists only of bounded operators. Hence

$$L^\infty(M) = \{h \in M_0 \mid \theta_s h = h, s \in \hat{\mathbb{R}}\} = M.$$

#### 1.9 Proposition

Let  $p \in [1, \infty[$  and let  $a$  be a closed, densely defined operator affiliated with  $M_0$ , and let  $a = u|a|$  be its polar decomposition. The following conditions are equivalent:

- (1)  $a \in L^p(M)$   
(2)  $u \in L^\infty(M)$  and  $|a|^p \in L^1(M)$ .

#### 1.10 Definition

On  $L^p(M)$  we define  $\|\cdot\|_p$  by

$$\|a\|_p = \text{tr}(|a|^p)^{\frac{1}{p}} \quad p < \infty$$
$$\|a\|_\infty = \|a\|$$

For  $p = 1, \infty$   $\|\cdot\|_p$  is a norm (cf. prop. 1.5). It will be proved later, that  $\|\cdot\|_p$  is also a norm for  $1 < p < \infty$ .

### 1.11 Lemma

Let  $h, k \in L^1(M)_+$ . The function  $\alpha \rightarrow h^\alpha k^{1-\alpha} \in L^1(M)$  is analytic in the open strip  $\{\alpha \in \mathbb{C} \mid \operatorname{Re} \alpha \in ]0, 1[ \}$ .

### 1.12 Proposition

Let  $p, q \in [1, \infty]$ ,  $\frac{1}{p} + \frac{1}{q} = 1$ , and let  $a \in L^p(M)$  and  $b \in L^q(M)$ , then  $ab, ba \in L^1(M)$  and  $\operatorname{tr}(ab) = \operatorname{tr}(ba)$ .

### 1.13 Corollary

(1) For any  $h \in L^1(M)$  and any unitary  $u \in L^\infty(M)$ :

$$\operatorname{tr}(uhu^*) = \operatorname{tr}(h)$$

(2) For any  $x \in L^2(M)$ :  $\operatorname{tr}(x^*x) = \operatorname{tr}(xx^*)$

### 1.14 Theorem

Let  $p, q \in [1, \infty]$ ,  $\frac{1}{p} + \frac{1}{q} = 1$ , and let  $a \in L^p(M)$  and  $b \in L^q(M)$ ,

then  $\|ab\|_1 \leq \|a\|_p \|b\|_q$  (Hölders inequality)

### 1.15 Remarks

The proof of Theorem 1.14 is based on lemma 1.11 and the three line theorem for analytic functions (compare with [9] p. 113). Dixmiers proof of Hölders inequality in [4] can not be applied, because in our case the spectral projections of an  $L^p$ -operator is not in  $L^p$ .

### 1.16 Proposition

(1) Let  $p, q \in [1, \infty]$ ,  $\frac{1}{p} + \frac{1}{q} = 1$ . For any  $a \in L^p(M)$

$$\|a\|_p = \sup \{ |\operatorname{tr}(ab)| \mid b \in L^q(M), \|b\|_q \leq 1 \}$$

(2) For  $a, b \in L^p(M)$   $\|a+b\|_p \leq \|a\|_p + \|b\|_p$ . (Minkowskis inequality)

Hence  $\|\cdot\|_p$  is a norm.

### 1.17 Proposition

(1) For any  $p \in [1, \infty]$  the norm topology on  $L^p(M)$  is equal to the topology of convergence in measure (cf [9] p. 106).

(2) For any  $p \in [1, \infty]$   $L^p(M)$  is complete in the  $p$ -norm.

(3)  $L^2(M)$  is a Hilbert space with inner product  $(a|b) = \operatorname{tr}(b^*a)$ .

1.18 Lemma (cf. [4] lemma 5 p.30)

Let  $p \in [2, \infty[$ . For  $a, b \in L^p(M)$

$$\|a + b\|_p^p + \|a - b\|_p^p \leq 2^{p-1}(\|a\|_p^p + \|b\|_p^p)$$

The proof of lemma 1.18 is based on lemma 1.11 and the three line theorem.

1.19 Theorem

Let  $p \in [1, \infty[$  and put  $q = \frac{p}{p-1}$ . An operator  $a \in L^q(M)$  defines a functional  $\varphi_a$  on  $L^p(M)$  by  $\varphi_a(x) = \text{tr}(ax)$ . The map  $a \rightarrow \varphi_a$  is an isometric isomorphism of  $L^q(M)$  onto the dual Banach space of  $L^p(M)$ .

Proof: Same as [4] proof of theorem 7.

1.20 Proposition

Let  $p, q \in [1, \infty]$ ,  $\frac{1}{p} + \frac{1}{q} = 1$  and let  $a \in L^q(M)$ . Then

$$a \geq 0 \iff \text{tr}(ab) \geq 0 \quad \forall b \in L^p(M)_+$$

i.e. the partial ordering of  $L^q(M)_{sa}$  is the dual of the ordering of  $L^p(M)_{sa}$ . ( $sa$  = selfadjoint).

For  $a \in M = L^\infty(M)$  and  $x \in L^2(M)$  we put

$$\begin{aligned} \lambda(a)x &= ax \\ \varrho(a)x &= xa. \end{aligned}$$

1.21 Theorem

- (1)  $\lambda$  (resp.  $\varrho$ ) is a normal, faithful representation (resp. anti-representation) of  $M$  on the Hilbert space  $L^2(M)$ .
- (2) The von Neumann algebras  $\lambda(M)$  and  $\varrho(M)$  are commutants of each other, and

$$\varrho(M) = J \lambda(M) J$$

where  $J$  is the conjugate linear isometry  $x \rightarrow x^*$  in  $L^2(M)$ .

- (3) The quadruple  $(\lambda(M), L^2(M), J, L^2(M)_+)$  is a standard form in the sense of [5].

## §2 The semifinite case

Let  $M$  be a semifinite von Neumann algebra on a Hilbert space  $H$ , and let  $\tau_0$  be a n.f.s. trace on  $M$ . Identifying  $L^2(\mathbb{R}, H)$  with  $H \otimes L^2(\mathbb{R})$  we have:

$$R(M, \sigma^{\tau_0}) = M \otimes U(\mathbb{R})$$

where  $U(\mathbb{R})$  is the von Neumann algebra associated with the left regular representation of the group  $\mathbb{R}$ . Let  $F$  denote the Fourier-Plancherel operator  $L^2(\mathbb{R}) \rightarrow L^2(\hat{\mathbb{R}})$

$$(Ff)(s) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{-ist} f(t) dt$$

$$\text{then } U(\mathbb{R}) = F^* L^\infty(\hat{\mathbb{R}}) F$$

where  $L^\infty(\hat{\mathbb{R}})$  acts as multiplication operators on  $L^2(\hat{\mathbb{R}})$ . Hence

$$R(M, \sigma^{\tau_0}) = M \otimes F^* L^\infty(\hat{\mathbb{R}}) F.$$

For any borel function  $f(s)$  on  $\hat{\mathbb{R}}$  we let  $m(f)$  denote the closed, densely defined multiplication operator  $g \rightarrow fg$  on  $L^2(\hat{\mathbb{R}})$ .

### 2.1 Theorem

Let  $p \in [1, \infty[$ . If  $a \in L^p(M, \tau_0)$  then  $a \otimes F^* m(\exp(\frac{\xi}{p})) F \in L^p(M)$  and the map  $a \rightarrow a \otimes F^* m(\exp(\frac{\xi}{p})) F$  is an isometry of  $L^p(M, \tau_0)$  onto  $L^p(M)$ .

## §3 Applications to von Neumann algebras with a periodic weight.

Let  $M$  be a von Neumann algebra with a periodic NSF-weight  $\varphi_0$ , and let  $T_0$  be a period for  $\varphi_0$ , i.e.  $\sigma_{T_0}^{\varphi_0} = 1$ . Put  $G = \mathbb{R}/T_0\mathbb{Z}$  and let  $t \rightarrow \dot{t}$  be the quotient map  $\mathbb{R} \rightarrow G$ . Let  $\alpha$  be the action  $\alpha: G \rightarrow \text{aut}(M)$  defined by  $\alpha(\dot{t}) = \sigma_t^{\varphi_0}$ . We will identify  $M$  with its injection in the crossed product  $M_1 = R(M, \alpha)$ .  $M_1$  is generated by  $M$  and a group of unitaries  $\lambda(g)$ ,  $g \in G$ , such that

$$\sigma_t^{\varphi_0}(x) = \lambda(\dot{t})x\lambda(\dot{t}) \quad x \in M.$$

Let  $S$  denote the operator valued weight  $M_1^+ \rightarrow \hat{M}_+$  given by

$$S(x) = \sum_{n=-\infty}^{\infty} \Theta^n(x) \quad x \in M_1^+$$

where  $\Theta^n$ ,  $n \in \mathbb{Z}$ , is the dual action of  $\hat{G} \simeq \mathbb{Z}$  on  $M_1$ . (c.f. [7]).

Repeating the arguments from the start of §1 we get that  $M_1$  has a (unique) n.f.s. trace  $\tau$  such that  $\varphi \circ S = \tau(k \cdot)$  where  $k$  is the positive selfadjoint operator affiliated with  $M_1$  determined by  $k^{it} = \lambda(i)$ ,  $t \in \mathbb{R}$ . The trace  $\tau$  satisfies:

$$\tau \circ \Theta = \lambda \tau \quad \text{where } \lambda = \exp(-\frac{2\pi}{i_0}).$$

### 3.1 Definition

For any normal semifinite weight  $\varphi$  on  $M$ , we put  $\tilde{\varphi} = \varphi \circ S$ , and we let  $k_\varphi$  be the Radon-Nikodym derivative of  $\tilde{\varphi}$  with respect to  $\tau$ , i.e.  $\tilde{\varphi} = \tau(k_\varphi \cdot)$ .

### 3.2 Proposition

(1) The set  $\{k_\varphi \mid \varphi \text{ normal, semifinite}\}$  is equal to the set of positive selfadjoint operators  $k$  affiliated with  $M_1$  for

$$\Theta k = \lambda k \quad (\lambda = \exp(-\frac{2\pi}{i_0})).$$

(2) Let  $k_\varphi = \int_0^\infty \mu de_\mu^\varphi$  be the spectral decomposition of  $k_\varphi$ , then for any  $a > 0$ :

$$\varphi(1) = \tau \left( \int_{\lambda a}^a \mu de_\mu^\varphi \right)$$

and

$$\frac{\lambda \varphi(i)}{(1-\lambda)a} \leq \tau((e_a^\varphi)^\perp) \leq \frac{\varphi(1)}{(1-\lambda)a}$$

in particular  $k_\varphi$  is  $\tau$ -measurable iff  $\varphi$  is bounded.

We could now continue as in §1 and construct new  $L^p$ -spaces consisting of the  $\tau$ -measurable operators affiliated with  $M_1$  which satisfies:

$$\begin{cases} \Theta k = \lambda^{\frac{1}{p}} k & p < \infty \\ \Theta k = k & p = \infty \end{cases}$$

However it is not hard to prove that these spaces are isomorphic to the spaces  $L^p(M)$  obtained from the general construction.

We will instead use proposition 3.2 to prove the following slight strengthening of a result due to Connes and Takesaki (cf. [3, Chap.II, corollary 4.10]).



### 3.3 Theorem

Let  $M$  be a  $\sigma$ -finite factor of type III  $\lambda, \lambda \in ]0, 1[$ .

- (1) For any two normal, faithful states  $\varphi, \psi$  on  $M$ , there exists a unitary  $u \in M$ , such that  $\lambda \varphi \leq u \varphi u^* \leq \lambda^{-1} \psi$ .
- (2) For any two unbounded n.f.s. weights  $\varphi, \psi$  on  $M$ , there exists a unitary  $u \in M$ , such that  $\lambda \varphi \leq u \varphi u^* \leq \lambda^{-1} \psi$ .

(The method of [3] gives  $\lambda^2 \varphi \leq u \varphi u^* \leq \lambda^{-2} \psi$  in the above inequalities).

### 3.4 Remark

It is easy to prove, that Theorem 3.3 is not valid for  $\lambda = 1$ .

A. Connes and E. Størmer [2] have recently proved, that any two normal states  $\varphi, \psi$  on a  $\sigma$ -finite type III<sub>1</sub>-factor are almost equivalent in the sense, that there exists a sequence of unitaries  $(u_n)_{n \in \mathbb{N}}$  in  $M$ , so that  $\|\psi - u_n \varphi u_n^*\| \rightarrow 0$  for  $n \rightarrow \infty$ .

### References

- [1] A. Connes, Une classification des facteurs de type III. Ann. Sc. de l'Ecole Normale Supérieure, 4 ser. t. 6, fasc. 2 (1973) 133-252.
- [2] A. Connes & E. Størmer, Homogeneity of the state space of factors of type III. J. funct. analysis 28 (1978) 187-196.
- [3] A. Connes & M. Takesaki, The flow of weights on a factor of type III. Tohoku Math. J. 29 (1977) 473-475.
- [4] J. Dixmier, Formes linéaires sur un anneau d'opérateurs. Bull. Soc. Math. France t. 81 (1953) 9-39.
- [5] U. Haagerup, The standard form of von Neumann algebras. Math. Scand. 37 (1975) 271-283.
- [6] U. Haagerup, Operator Valued Weights in von Neumann algebras. To appear in J. funct. analysis.

- [7] U. Haagerup , On the dual weights for crossed products of von Neumann algebras II. Math. Scand. 43,119-140 (1978).
- [8] R. Kunze ,  $L_p$  Fourier transform on locally compact unimodular groups. Trans. Amer. Math. Soc. 89 (1958) 519-540.
- [9] E. Nelson , Notes on non-commutative integration. Journal of funct. anal. 15 (1974) 103-116.
- [10] G.K. Pedersen & M. Takesaki , The Radon Nikodym Theorem for von Neumann Algebras. Acta Math. 130 (1973) 53-87.
- [11] I. Segal , A non commutative extension of abstract integration. Ann. of Math. 57 (1953) 401-457.
- [12] M. Takesaki , Duality for crossed products and the structure of von Neumann algebras of type III. Acta Math. 131, (1973) 249-308.

Uffe Haagerup

Matematisk Institut, Odense Universiteit

Niels Bohrs Allé, 5000 Odense, Denmark.